



FALLS PREVENTION SOLUTION

YELLOW SLIPPER SOCKS FOR HIGH-RISK PATIENTS

Our Recommendations

In accordance with this trust's falls prevention policy, we recommend that Medline Slipper Socks be used as a method of reducing slips and trips that can lead to falls in the following situations:

- Patients have no, or inappropriate, footwear on admission to the hospital or whilst in your department.
- Regular footwear cannot be applied (e.g. over dressings or oedematous feet).
- A patient is non-concordant in wearing footwear at all the recommended times (e.g. due to confusion or agitation) as being barefoot or wearing conventional socks puts the patient at a greater falls risk.

The slipper socks **MUST BE** removed at least daily to review foot health and hygiene and to perform pressure area checks.

Caution should be taken when applying slipper socks to patients with:

- Unstable gait
- Lower limb vascular disease
- Lower limb compression dressings
- Lower limb oedema
- Foot disorders
- Diabetic neuropathy.

Please refer to the appropriate specialist if you have questions about whether slipper socks are an appropriate intervention for your patient.

Medline SKU	NPC Code	Sock Size*	UK Size	Colour	Packaging
MDTES4B04FPY	CVW050	Large	3-5		48 pr/cs
MDTES4B05FPY	CVW051	X-Large	5-9		
MDTES4B06FPY	CVW052	XX-Large	9-11		

*We also offer Child and Medium sock sizes, but Large to XX-Large covers the vast majority of patient needs.

For further information, please refer to your Medline Falls Prevention Pocket Guide or contact [insert name of appropriate person(s)].

sales rep details

